

# Physics potentials of a magnetized iron calorimeter detector

Srubabati Goswami

Physik-Department T30d  
Technische Universität München , Germany

&

Harish-Chandra Research Institute,  
Allahabad, India

**For the INO collaboration**

**SNOW2006**

# Magnetized Iron Calorimeter Detector



---

- Currently feasibility study for such a detector is underway in India by the **India-Based Neutrino Observatory (INO)** collaboration.
- Detector choice based on
  - Technological capabilities available in the country
  - Existing/Planned other neutrino detectors in the world
  - Modularity and the possibility of phasing
  - Compactness and ease of construction
- **MONOLITH** collaboration had earlier proposed similar design

# The detector

- Magnetised iron calorimeter ( $\sim 50\text{kT}$ )
- 140 horizontal (vertical) iron layers interspersed with Glass RPC
- Modular structure

- Sensitive to muons
- Energy determination from
  - Track length
  - Track curvature in a magnetic field
- Direction of parent neutrino from the track
- Charge identification from track curvature in magnetic field



# Current Activities

---


- Detector R & D
- Physics Studies
- Detector Simulation
- Data Acquisition System
- Site Survey
- Human Resource Development

Interim Report submitted to funding agencies

# Cost Estimates and Time Schedule

---

## Cost

-  Lab. Construction  $\sim$  90 crores INR (1 crore = 10 million)
-  Detector  $\sim$  (200 (iron) + 200 (others)) crores in INR


Total cost  $\sim$  500 crores in INR (1 Euro  $\approx$  INR 50)

## Time Scale : $\sim$ 5 years from approval

Details: INO interim report, <http://www.imsc.res.in/~ino>

# Site

- Two sites were considered – **Rammam** in North India and **PUSHEP** in South India
- PUSHEP is recommended for ease of accessibility, less seismicity..




Geotechnological studies are going on

# Physics Goals for INO


---

- First phase – measurement of atmospheric neutrino flux
  - Reconfirmation of the first oscillation dip as a function of L/E
  - Improved precision of oscillation parameters
  - Determination of the octant of  $\theta_{23}$
  - Matter effects and determination of sign of  $\Delta m_{31}^2$
  - Probing CPT violation, Lorentz violation
  - Discrimination between  $\nu_{\mu} - \nu_{\tau}$  and  $\nu_{\mu} - \nu_s$
  - Constraining long range leptonic forces
- Second Phase – end detector for beta beams, neutrino factory
  - hierarchy,  $\theta_{13}$ , CP violation
  - CERN to INO baseline  $\sim 7000$  km, the magic baseline

# INO as a long baseline detector





# Disappearance of $\nu_\mu$ vs $L/E$



$$\frac{N_{up}(L/E)}{N_{down}(L/E)} \simeq P_{\mu\mu}$$


$$= 1 - \sin^2 2\theta_{23} \sin^2 \Delta_{31} L/4E$$

Expect to determine  $\Delta_{31}$  with 10% precision




# Ambiguity in Mass Hierarchy

$$\tan 2\theta_{13}^m = \frac{\Delta m_{31}^2 \sin 2\theta_{13}}{\Delta m_{31}^2 \cos 2\theta_{13} \pm 2\sqrt{2}G_F n_e E}$$



- For  $\Delta m_{\text{atm}}^2 > 0$  matter resonance in neutrinos
- For  $\Delta m_{\text{atm}}^2 < 0$  matter resonance in anti neutrinos
- Experiments sensitive to **matter effects** can probe the mass hierarchy
- Matter effects** for  $\Delta m_{\text{atm}}^2$  channel depend crucially on  $\theta_{13}$
- Thus both parameters get related

# Ambiguity in Mass Hierarchy




M. Lindner, hep-ph/0503101

- Hierarchy difficult to determine in superbeams
- Sensitivity limited by correlation and degeneracies
- Synergistic use of experiments
- Use of Magic Baseline

# Earth Matter Effects at Long Baselines

Problem of  $\delta_{CP}$  degeneracy less at longer baselines



Significant matter effect in  $P_{\mu\tau}$  at 9700 km and for  $E \sim 5$  GeV

Genuine three flavour effect

Impact on  $P_{\mu\mu} \Rightarrow$

$$P_{\mu\mu} = 1 - P_{\mu e} - P_{\mu\tau}$$

At 7000 km drop in  $P_{\mu\mu}$  induced by  $P_{\mu e}$

At 9700 km rise in  $P_{\mu\mu}$  induced by  $P_{\mu e}$  and  $P_{\mu\tau}$

R. Gandhi et. al, PRL, 2005

# Determining Hierarchy by Atmospheric Neutrinos

- Using  $\mu^-$  rates in magnetized iron calorimeter detectors like **INO**

$$\begin{aligned}\phi_{\mu^-} / \phi_{\mu^-}^0 &\approx P_{\mu\mu} + rP_{e\mu} & r &= \phi_e^0 / \phi_{\mu}^0 \\ &= P_{\mu\mu}(1 - r) - rP_{\mu\tau} + r\end{aligned}$$



- For  $\Delta m_{31}^2 > 0$  matter effect in  $\nu_{\mu}$  ( $N_{\mu+}^{\text{mat}} \approx N_{\mu+}^{\text{vac}}$ )

# Determining Hierarchy by Atmospheric Neutrinos

Using  $\mu^-$  rates in magnetized iron calorimeter detectors like INO

$$\begin{aligned} \phi_{\mu^-} / \phi_{\mu^-}^0 &\approx P_{\mu\mu} + rP_{e\mu} & r &= \phi_e^0 / \phi_{\mu}^0 \\ &= P_{\mu\mu}(1 - r) - rP_{\mu\tau} + r \end{aligned}$$

For  $\Delta m_{31}^2 > 0$  matter effect in  $\nu_{\mu}$  ( $N_{\mu^+}^{\text{mat}} \approx N_{\mu^+}^{\text{vac}}$ )



Gandhi et al., hep-ph/0411252

Palomarez-Ruiz, hep-ph/0406096

Murthy, Indumathi hep-ph/0407336

# Determining Hierarchy by Atmospheric Neutrinos

- Using  $\mu^-$  rates in magnetized iron calorimeter detectors like **INO**

$$\begin{aligned}\phi_{\mu^-} / \phi_{\mu^-}^0 &\approx P_{\mu\mu} + rP_{e\mu} & r &= \phi_e^0 / \phi_{\mu}^0 \\ &= P_{\mu\mu}(1 - r) - rP_{\mu\tau} + r\end{aligned}$$

- For  $\Delta m_{31}^2 > 0$  matter effect in  $\nu_{\mu}$  ( $N_{\mu^+}^{\text{mat}} \approx N_{\mu^+}^{\text{vac}}$ )
- 3-4 $\sigma$  signal for matter effects at  $\sin^2 2\theta_{13} = 0.1$  for 1000kTy using the total event rates **for fixed values of parameters**
- Parameter uncertainties spoil the sensitivity

# Bin by bin $\chi^2$ -analysis

## Results for a iron calorimeter detector

- $\chi^2$  analysis of  $\mu^-$  event in 24 L/E bins
- 15% energy and  $15^\circ$  angular resolution
- 10% systematic error
- 85% efficiency
- Marginalized over  $\Delta m_{31}^2, \sin^2 \theta_{13}, \sin^2 \theta_{23}$

$\sin^2 2\theta_{13}$	$\chi_{\min}^2$ 500 kt yr	$\chi_{\min}^2$ 1000 kt yr
0.05	2.7	3.7
0.1	6.6	8.9

Gandhi et al. work in progress.

# Bin by bin $\chi^2$ -analysis


## Results for a iron calorimeter detector

- $\chi^2$  analysis of  $\mu^-$  event in 24 L/E bins
- 15% energy and 15° angular resolution
- 10% systematic error
- 85% efficiency
- Marginalized over  $\Delta m_{31}^2$ ,  $\sin^2 \theta_{13}$ ,  $\sin^2 \theta_{23}$

$\sin^2 2\theta_{13}$	$\chi_{\min}^2$ 500 kt yr	$\chi_{\min}^2$ 1000 kt yr
0.05	2.7	3.7
0.1	6.6	8.9

Gandhi et al. work in progress.

## Effect of Smearing



Petcov and Schwetz, hep-ph/0511277

# Bin by bin $\chi^2$ -analysis

## Results for a iron calorimeter detector

- $\chi^2$  analysis of  $\mu^-$  event in 24 L/E bins
- 15% energy and  $15^\circ$  angular resolution
- 10% systematic error
- 85% efficiency
- Marginalized over  $\Delta m_{31}^2, \sin^2 \theta_{13}, \sin^2 \theta_{23}$

$\sin^2 2\theta_{13}$	$\chi_{\min}^2$ 500 kt yr	$\chi_{\min}^2$ 1000 kt yr
0.05	2.7	3.7
0.1	6.6	8.9

Gandhi et al. work in progress.

## Comparison with water-Cerenkov detector

- No charge sensitivity:  $N_\mu = N_\mu^+ + N_\mu^-$

$\sin^2 2\theta_{13}$	$\chi_{\min}^2$ (6 Mt yr)
0.05	1.9
0.1	4.4


Gandhi et al., hep-ph/0406145

# Deviation of $\sin^2 \theta_{23}$ from maximal value

---

- $D \equiv 1/2 - \sin^2 \theta_{23}$
- $|D|$  gives the deviation of  $\sin^2 \theta_{23}$
- $\text{sgn}(D)$  gives the octant of  $\sin^2 \theta_{23}$
- Current  $3\sigma$  limits:
  - $|D| < 0.16$  at  $3\sigma$  from the SK data
  - No robust information on  $\text{sgn}(D)$

# Can Earth matter effects determine $|D|$ ?



$$P_{\mu\mu}^m = 1 - P_{\mu\mu}^{m1} - P_{\mu\mu}^{m2} - P_{\mu\mu}^{m3}$$

$$P_{\mu\mu}^{m1} = c_{13}^2 \sin^2 2\theta_{23} \sin^2 [1.27(\Delta_{31} + A + \Delta_{31}^m)L/2E]$$

$$P_{\mu\mu}^{m2} = s_{13}^2 \sin^2 2\theta_{23} \sin^2 [1.27(\Delta_{31} + A - \Delta_{31}^m)L/2E]$$

$$P_{\mu\mu}^{m3} = \sin^4 \theta_{23} \sin^2 2\theta_{13}^m \sin^2 (1.27\Delta_{31}^m L/E)$$



Dependence on  $\theta_{23}$  in the form  $\sin^4 \theta_{23}$



Octant sensitivity ?


S.Choubey. and P. Roy hep-ph/0509197

Also Indumathi et al. hep-ph/0603264

# Can Earth matter effects determine $|D|$ ?



Using atmospheric neutrinos in INO




$|D|$  can be measured to  $\sim 17\%$  ( $20\%$ ) at  $3\sigma$  for  $s_{13}^2 = 0.04$  ( $0.00$ ) with 1 MtonY exposure and 50% detector efficiency

S.Choubey. and P. Roy hep-ph/0509197

# Resolving the octant ambiguity in INO

- Using atmospheric neutrinos in INO
- For every non-maximal  $\sin^2 \theta_{23}(\text{true})$  there exists a  $\sin^2 \theta_{23}(\text{false})$   
$$\sin^2 \theta_{23}(\text{false}) = 1 - \sin^2 \theta_{23}(\text{true})$$



S.Choubey. and P. Roy hep-ph/0509197

# Comparing the Octant Sensitivity of Experiments

## Long baseline experiments

No octant sensitivity

 LBL+atmospheric Huber et al hep-ph/0501037

 LBL accelerator + reactor Minakata et al hep-ph/0601258

## Atmospheric neutrinos in water Cerenkov detectors

$\sin^2 \theta_{23}(\text{false})$  can be excluded at  $3\sigma$  if:

$$\sin^2 \theta_{23}(\text{true}) < 0.36 \text{ or } > 0.62$$

Gonzalez-Garcia et al, hep-ph/0408170

## Atmospheric neutrinos in large magnetized iron detectors

$\sin^2 \theta_{23}(\text{false})$  can be excluded at  $3\sigma$  if:

$$\sin^2 \theta_{23}(\text{true}) < 0.36 \text{ or } > 0.63 \text{ for } \sin^2 \theta_{13}(\text{true}) = 0.01,$$

$$\sin^2 \theta_{23}(\text{true}) < 0.40 \text{ or } > 0.59 \text{ for } \sin^2 \theta_{13}(\text{true}) = 0.02,$$

$$\sin^2 \theta_{23}(\text{true}) < 0.41 \text{ or } > 0.58 \text{ for } \sin^2 \theta_{13}(\text{true}) = 0.03,$$

$$\sin^2 \theta_{23}(\text{true}) < 0.42 \text{ or } > 0.57 \text{ for } \sin^2 \theta_{13}(\text{true}) = 0.04.$$

S.Choubey. and P. Roy hep-ph/0509197

# Detector and Physics Simulation

---

- Simulation studies with atmospheric neutrinos are in progress at many collaborating Institutions

-  **Nuance Event Generator**

-  Generates of atmospheric neutrino events inside the INO detector

-  **GEANT Monte Carlo Package**

-  Simulates the detector response for the neutrino events

-  **Event Reconstruction**

-  Fits the raw data to extract neutrino energy and direction

-  **Physics Performance**

-  Analysis of reconstructed events to extract physics.

# Conclusion

---

- A large magnetized iron calorimeter detector has substantial physics potential using atmospheric neutrinos.
  - Reconfirmation of L/E dip and precision of  $\Delta m_{31}^2$
  - Matter effect and Sign of  $\Delta m_{31}^2$
  - Determination of octant of  $\theta_{23}$
  - CPT violation, Long Range Forces .....
- It will complement the planned water Cerenkov, Liquid Scintillator and Liquid Argon Detectors as well as the long baseline and reactor experiments
- Can be used as a far detector for neutrino factories

Should be an International Facility